

Shotgun Femtosecond Diffractive Imaging of Free Nanoscale Biomaterials

Michael J. Bogan¹, Urs Rohner¹, Sébastien Boutet^{1,2,3}, W. Henry Benner¹, Anton Barty¹, Saša Bajt¹, Matthias Frank¹, Bruce Woods¹, Stefano Marchesini^{1,4}, Stefan P. Hau-Riege¹, Marvin Seibert³, Filipe Maia³, Florian Burmeister³, Erik Marklund³, Rolf Treusch⁵, Eberhard Spiller⁶, Thomas Möller⁷, Christoph Bostedt⁷, Janos Hajdu^{2,3}, and Henry N. Chapman^{1,4}

1. University of California, Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore CA 94550, USA. 2. Stanford Synchrotron Radiation Laboratory, Stanford Linear Accelerator Center, 2575 Sand Hill Road, Menlo Park, California 94305, USA. 3. Laboratory of Molecular Biophysics, Institute of Cell and Molecular Biology, Uppsala University, Husargatan 3, Box 596, S-75124 Uppsala, Sweden 4. Center for Biophotonics Science and Technology, University of California, Davis, 2700 Stockton Blvd., Suite 1400, Sacramento, CA 95817, USA. 5. Deutsches Elektronen-Synchrotron, DESY, Notkestraße 85, D-22607 Hamburg, Germany 6. Spiller X-ray Optics, Livermore CA 94550, USA. 7. Institut für Atomare Physik, Technische Universität Berlin, Hardenbergstraße 36, PN 3-1, 10623 Berlin, Germany

Femtosecond diffractive imaging is a new x-ray microscopy technique for imaging non-crystalline objects beyond the damage threshold. Part of a new era of X-ray science delivered by X-ray free electron lasers (FEL), this technique has the potential to deliver three-dimensional atomic resolution structures of non-crystalline nanoscale biomaterials such as single biomolecules using images collected from reproducible copies exposed to the beam one by one. Experimentally, container-less delivery of single biomolecules to the X-ray pulses is imperative because any atom present in the X-ray path will contribute to the diffraction pattern. Here we use the soft-X-ray FEL in Hamburg (FLASH) to perform the first demonstration femtosecond diffractive imaging of free nanoscale biomaterials via a shotgun approach—a continuously refreshed stream of single nanoparticles synthesized *in situ* at atmospheric pressure by charge-reduced electrospray of a sucrose solution containing megadalton DNA origami complexes. In our method, the aerosol is transformed into a tightly focused particle stream in-vacuum using a set of aerodynamic lenses and single events of the interception of individual nanoparticles with an intense 10 femtosecond pulse, containing $\sim 10^{12}$ photons at 13.5 nm, results in a coherent diffraction pattern captured with a single photon sensitive X-ray camera. Reconstructed images of intercepted particles are obtained by phase retrieval through oversampling. Ions generated from the explosion of the nanoparticle are detected by a miniature mass spectrometer, providing insights to chemical composition. In its current configuration shotgun femtosecond diffractive imaging operates at 0.1 Hz and is directly transferable to future hard-X-ray FEL sources.